



THE KENYA VETERINARY VACCINES PRODUCTION INSTITUTE

FOWLVAX™

Inactivated *Salmonella gallinarum* (Fowl typhoid) Vaccine.

COMPOSITION

FOWLVAX™ is a multi-strain inactivated Fowl typhoid vaccine prepared from *Salmonella gallinarum*. The strains included are CN 175, 176, 177, 178, and 179.

PROPERTIES

Fowl typhoid is a disease of chickens caused by *Salmonella gallinarum* and is more often observed in the later growing period and in mature stock. The clinical signs resemble those of septicaemia, with inappetence and, eventually, death.

INDICATIONS

For the active immunization of healthy chicken against Fowl typhoid.

DOSAGE AND ADMINISTRATION

Shake well before use. Vaccination should be done using sterile needle and syringe to avoid contamination. Administer to birds via intramuscular injection in the thigh or breast muscles at the rate of 1ml per bird.

CONTRA-INDICATIONS, WARNINGS

For Veterinary use only. Do not use if seal is broken or package is damaged. Do not vaccinate un-healthy birds. This preparation is strictly for use in poultry only. All vaccines may occasionally cause anaphylactic reactions in some birds. If this occurs adrenaline should be administered.

STORAGE

Store at +4°C. (Refrigerator). DO NOT FREEZE. If stored at between +2°C and +8°C (refrigerator) the shelf life is one year. Once the vaccine bottle has been broached it must be used immediately and any remaining quantity discarded.

PRESENTATION

50-, 100-, and 300-dose vials.